Mouse Anti-CA 125 Monoclonal Antibody Datasheet

Product Name: mAb anti-CA 125 Clone No.: 23

Catalogue No.: MO-T40012C Quantity: 0.5 mg/vial

Description: Mouse monoclonal antibody to human

cancer antigen 125 (CA125)

Purification: Protein G affinity purified

Product Type: Primary antibody

Target Protein: Human CA 125

Immunogen: Human CA 125 protein

Fusion Sp2/0-Ag14

Myeloma:

Specificity: React with human CA 125

Reactivity: Human, others not tested

Species Not available

Reactivity:

Host / Isotype: Mouse, IgG2a Kappa

Formulation: Lyophilized from a solution in 0.01M

PBS, pH 7.2

Reconstitution: Double distilled water is recommended

to reconstitute the antibody.

Storage: Store at -20°C

Research Oncology


Area:

Background: Cancer antigen 125 (CA125), also called

mucin16, is a heterogeneous mixture of glycoproteins with a molecular weight range of 200-1000 kDa. Its C-end is attached to cell membrane. The extracellular region of the molecule can be


released from cell surface by proteolytic cleavage. CA125 was found to be elevated in the blood of patients with advanced ovarian cancer. It is very useful for monitoring the effects of ovarian cancer therapy and is used for predicting the outcome of the patients after treatment.

Applications: 1. Fluorescence flow cytometry



Above is the histogram of Hela cells stained with $10\mu g/mL$ anti-CA 125 mAb clone 23 and fluorescence-labelled secondary antibody. Black line represents the histogram of Hela cells stained with control antibody, mouse anti-TB 38Kd antigen mAb clone B12F8 (MO-I40021D).

2. Immunofluorescence



This product is for LABORATORY RESEARCH USE and further manufacture ONLY, and cannot be administrated to human and animals for use in diagnostic and therapeutic procedures.

Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.

Page 1 of 2 page(s)

S7.5 (02)



ANOGEN - A Division of YES Biotech Laboratories Ltd.

2355 Derry Road East, Unit 23, Mississauga, ON, Canada L5S 1V6 • Tel: (905) 677-9221 • Fax: (905) 677-0023

Staining of methanol-fixed Hela cells by Immuno-fluorescence: Anti-CA 125 mAb clone 23 used at $10\mu g/mL$.

3. ELISA: This antibody, when used at 0.5ng/mL or higher concentrations, can recognize ELISA plate wells coated with CA125 at 5 international unit per well.

References:

If research is published using this product, please inform Anogen in order to cite the reference on this datasheet. Anogen will provide one unit of product in the same category as gratitude.